

## 最近の研究業績

- Nemoto K, Mihara K, Nakamura A et al: Effects of escitalopram on plasma concentrations of aripiprazole and its active metabolite, dehydroaripiprazole, in Japanese patients. *Pharmacopsychiatry*. 2014; 47: 101-4.
- Suzuki T, Mihara K, Nakamura A et al: Effects of genetic polymorphisms of *CYP2D6*, *CYP3A5*, and *ABCB1* on the steady-state plasma concentrations of aripiprazole and its active metabolite, dehydroaripiprazole, in Japanese patients with schizophrenia. *The Drug Monit*. 2014; 36: 651-5.
- Kagawa S, Mihara K, Nakamura A et al: Relationship between plasma concentrations of lamotrigine and its early therapeutic effect of lamotrigine augmentation therapy in treatment-resistant depressive disorder. *The Drug Monit*. 2014; 36: 730-3.
- Nakamura A, Mihara K, Nemoto K et al: Lack of correlation between the steady-state plasma concentrations of aripiprazole and haloperidol in Japanese patients with schizophrenia. *The Drug Monit*. 2014; 36: 815-8.
- 三原一雄、甲田宗良、中村明文ら: 気質が気分障害の臨床的特徴と治療反応性に与える影響－循環気質と発揚気質に焦点を当てて－. *臨床精神薬理*. 2014; 17: 175-80.
- 中村明文、三原一雄、香川祥子ら: Lamotrigine の双極性障害への適応. *臨床精神薬理*. 2014; 17: 971-8.